CAUTION

KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS FOR ANIMAL TREATMENT ONLY



advantage®

for Dogs & Cats

100 g/L IMIDACLOPRID

For the treatment and prevention of fleas (Ctenocephalides spp.) on dogs, cats, rabbits and ferrets. Larval flea stages are killed in the surroundings of ADVANTAGE® treated pets. Reduces the incidence of Flea Allergy Dermatitis. For the control of lice (Trichodectes canis, Linognathus setosus) on dogs. One application on the skin lasts for a month on dogs and up to a month on cats.

RESTRAINT: DO NOT USE in rabbits that may be consumed or processed for

For external application only

DOSAGE

For animals of this type \P		Use 1 TUBE from this pack colour ↓
Small cats & kittens	up to 4 kg	dark orange
Cats	over 4 kg	violet
Small dogs & puppies, rabbits and ferrets	up to 4 kg	dark green
Dogs, rabbits and ferrets	4-10 kg	turquoise
Dogs	10-25 kg	maroon
Dogs	25-50 kg	blue

For very large dogs over 50 kg, apply 2 tubes of ADVANTAGE® for Dogs over 25 kg (blue) using one tube between the shoulder blades and one tube on the midline of the back between the hips.

Remove required tube(s) from the package



- 2. Hold tube upright and pull off the cap
- $\beta. \;\;$ Reverse the cap and push cap back on tube.
- 4. Twist cap to break seal, then remove cap from tube.
- Site of Application

CATS	DOGS UNDER 25 kg	DOGS 25-50 kg		
1 TUBE Apply on the back	1 TUBE Apply between	1 TUBE Apply by dividing the		
of the neck.	the shoulder blades.	dose between the shoulder blades, on		
		the midline of the back		
		between the hips, and		
The unity		at one point in between.		

- The animal should be standing for easy application. Remove any collar prior to application and replace when product is dry. Part the hair until the skin is visible. Place the tip of the tube on the skin and squeeze the tube firmly twice to apply the solution directly on the skin. Avoid pet's eyes or mouth. If licked by cats, signs of distaste may be observed due
- Record treatment date and repeat monthly or as required, ADVANTAGE® remains efficacious following shampoo treatment, swimming or exposure to rain. Avoid applying the product while the animal is wet. If washing animal during use of this product use a soap free shampoo.
- 8. Occasionally, hair loss may occur at the application site. This is usually a single occurrence and hair will grow back uneventfully.

GENERAL INSTRUCTIONS:

ADVANTAGE® has been shown to kill adult and larval fleas within 20 minutes of contact by absorption via the flea intersegmental membranes. Stops fleas from feeding in 3-5 minutes. Fleas are not required to bite the animal to receive a lethal dose.

The successive feeding activity of fleas on pets may elicit a hypersensitivity skin disorder known as flea allergy dermatitis (FAD). Treatment of pets with ADVANTAGE® rapidly kills fleas and reduces the incidence of this condition

ADVANTAGE® is safe for use on pregnant and lactating animals.

It is not necessary to treat unweaned puppies and kittens as treatment of the mother protects the litter while in contact. Treat the mother until weaning, then treat individual puppies and kittens.

ADVANTAGE® may be used at the same time as treatment for worms with DRONTAL®/DRONCIT® and with daily or monthly preventive treatment

It is recommended that all dogs and cats in the same environment should be protected by ADVANTAGE $^{\otimes}.$

ADVANTAGE® may be used to prevent infestation in flea-free animals taken temporarily to known flea infested environments. Apply a single dose before moving to the infested environment.

ADVANTAGE® may be used as the only product for flea control as it breaks the life cycle and kills fleas emerging from the environment to reinfest the cat/dog. More rapid resolution of heavy environmental infestations may be obtained by additional environmental control measures using a suitable registered product. If re-treatment becomes necessary earlier than 4 weeks, do not re-treat more than once weekly.

Re-treatment of rabbits and ferrets may be necessary when flea infestation is heavy.

WARNING:

Take care when treating ferrets to administer the correct dose. Safety data on overdosing is not available.

ADVANTAGE® has no strong chemical smell. The liquid vehicle in ADVANTAGE® may discolour or damage certain surfaces and fabrics, e.g., polished floorboards, leather. Exclude the animal from contact with such surfaces until the product dries on the animal's coat.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

SAFETY DIRECTIONS:

Harmful if inhaled or swallowed. Will irritate the eyes. Avoid contact with eyes. Do not inhale vapour. If product in eyes, wash it out immediately with water. Wash hands after use.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

DISPOSAL:

Dispose of used tubes and packaging by wrapping in paper and placing

in garbage. PRESENTATION:

ADVANTAGE® for	Pack Colour	Contents	APVMA Approval No:
Kittens & Small Cats up to 4 kg	dark orange	0.4 mL	50395/57506
Cats over 4 kg	violet	o.8 mL	50396/57508
Puppies & Small Dogs up to 4 kg	dark green	0.4 mL	50397/57510
Dogs 4–10 kg	turquoise	1.0 mL	50398/57511
Dogs 10-25 kg	maroon	2.5 mL	50399/57513
Dogs over 25 kg	blue	4.0 mL	50400/57503

Store below 30°C (Room Temperature).

LIMITATION OF LIABILITY:

Failure to use the product strictly as directed may be illegal, prove dangerous or render the product ineffective. Notwithstanding anything else contained on this label, and subject to any rights you have under the Australian Consumer Law and any other applicable law, Bayer Australia Limited's liability is limited to replacing this product or supplying equivalent product or paying the cost of replacing this product, at its discretion

ADVANTAGE®, DRONTAL®, and DRONCIT® are registered trademarks of Bayer AG, Leverkusen, Germany.

BAYER AUSTRALIA LIMITED

875 Pacific Highway Pymble NSW 2073 ACN 000 138 714

Bayer

82113088 6/14