

FOR ANIMAL TREATMENT ONLY



DRONTAL®

Cat Allwormer and Large Cat Allwormer

DRONTAL CAT ALLWORMER
20 mg PRAZIQUANTEL, 80 mg PYRANTEL as the EMBONATE

DRONTAL LARGE CAT ALLWORMER
30 mg PRAZIQUANTEL, 120 mg PYRANTEL as the EMBONATE

Controls gastrointestinal worms and tapeworms.
Drontal is for the control of all gastrointestinal worms and tapeworms in cats including:

Roundworm *Toxocara cati*, *Toxascaris leonina*.
Hookworm *Ancylostoma tubaeforme*, *Ancylostoma braziliense* (this hookworm is rare outside the coastal areas of north Queensland, Northern Territory and northern Western Australia), *Uncinaria stenocephala*.
Tapeworm *Dipylidium caninum* (common flea tapeworm)
Taenia taeniaeformis

Does not control lungworm

DIRECTIONS FOR USE

Drontal Allwormer is given as a single dose by mouth.

Cats should be dosed on an empty stomach. It is essential to weigh cats before treatment.

DOSAGE

Bodyweight	Drontal Cat Allwormer Tablets	Drontal Large Cat Allwormer Tablets
2kg and under	1/2	
2.1-4kg	1	
4kg - 6kg	One tablet per 4kg	1
Over 6kg	One tablet per 4kg	

If worm problems persist consult a veterinarian

Worms, if present, may not appear in the droppings after treatment, or may not appear for up to 48 hours after treatment.

RECOMMENDED TREATMENTS

Worms often reinfest cats so it is necessary to:

- treat the cat itself,
- reduce environmental contamination by daily disposal of droppings where possible,
- control intermediate hosts eg. fleas, rats, mice.

This table summarises retreatment recommendations:

Kittens	Adults	Pregnant/Lactating Queens
Treat at 6, 8, 12 weeks; then 3 monthly.	Every 3 months.	Prior to mating, 10 days before kitting and 2 and 4 weeks after kitting.

These recommendations are of a general nature. In specific instances where infestation is particularly heavy or many cats are housed together, a veterinary surgeon should be consulted.

Flea tapeworms can reinfest cats and begin producing segments 2-3 weeks after a treatment. If segments reappear before the next DRONTAL treatment is due, either retreat with DRONTAL or use DRONCIT Tapewormer for Dogs and Cats as an intermediate, tapeworm only, treatment.

COMPATIBILITY

DRONTAL may be combined with other treatments, for example flea or tick treatment.

SAFETY

DRONTAL offers gentle, effective control of worms. It may be given to all cats including old animals, young kittens, pregnant queens and show animals.

This prescription is not formulated for dogs and is not suitable for this use.

NOTE

Zipper worm (*Spirometra erinacei*) is a less common tapeworm. If segments are seen in the droppings less than 2 weeks after treatment they will probably belong to this worm. When tapeworm treatment only is required, as in this case, give DRONCIT Tapewormer for Dogs and Cats which is specifically for tapeworm treatment.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

DISPOSAL

Dispose of empty container by wrapping with paper and putting in garbage.

STORAGE

Store below 30°C (Room Temperature).

LIMITATION OF LIABILITY:

Failure to use the product strictly as directed may be illegal, prove dangerous or render the product ineffective. Notwithstanding anything else contained on this label, and subject to any rights you have under the Australian Consumer Law and any other applicable law, Bayer Australia Limited's liability is limited to replacing this product or supplying equivalent product or paying the cost of replacing this product, at its discretion.

DRONTAL® and DRONTAL ALLWORMER® are registered trademarks of Bayer AG, Leverkusen, Germany.

Drontal for	APVMA Approval No.
Cats up to 4 kg	46718/59170
Cats over 4 kg	53751/59169

BAYER AUSTRALIA LIMITED
875 Pacific Highway Pymble NSW 2073
ACN 000 138 714

Bayer

XXXXXX_05/14