

PRINT SIDE

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY



kiltix®

for dogs

Tick Collar

ACTIVE CONSTITUENTS: 22.5g/kg FLUMETHRIN, 100g/kg PROPOXUR (AN ANTICHOLINESTERASE COMPOUND)

- Aids in control of paralysis ticks (*Ixodes holocyclus*) for 6 weeks
- Controls brown dog and bush ticks for up to 5 months
- Controls fleas for 5 months
- Adjustable to fit all dog sizes



kiltix®

Tick Collar for dogs

- Aids in control of paralysis ticks (*Ixodes holocyclus*) for 6 weeks
- Controls brown dog and bush ticks for up to 5 months
- Controls fleas for 5 months
- Adjustable to fit all dog sizes

Batch:
Expiry:

kiltix®

Tick Collar for dogs

For a free treatment reminder visit
www.friendsforever.com.au

kiltix®

Tick Collar for dogs

- Control of ticks (*Rhipicephalus sanguineus*, *Haemaphysalis longicornis*) for up to 5 months.
- Aids in control of Paralysis Ticks (*Ixodes holocyclus*) for 6 weeks.
- Controls fleas (*Ctenocephalides* spp.) for 5 months.
- Collar length 66cm.

DIRECTIONS FOR USE

This product is contraindicated for use on sick or convalescing dogs or on puppies less than 3 months old.

This product is contraindicated for use with any other anticholinesterase pesticide on the dog while the collar is worn. Do not allow pets to chew on the collar.

Remove the collar from the pouch. Unroll the collar, place around the dog's neck and pull the free end through the clasp. Leave space enough for two fingers under collar. For smaller dogs cut the extra collar length off and dispose of cut off end wrapped in paper. The collar should be worn continuously.

The safety of KILTIX has not been established during pregnancy and lactation. However evidence suggests that no adverse effects are to be expected in pregnant and lactating bitches.

GENERAL INSTRUCTIONS

The period of control will vary with environmental conditions so the collar may need to be replaced at more frequent intervals. While fleas and ticks are present, the collar should be worn constantly. Replace when effectiveness diminishes. When this collar is first worn, observe the neck area every few days for signs of irritation as some dogs may be sensitive to collars. Remove the collar at the first sign of irritation or adverse reaction. This collar resists the effect of water, nevertheless it is recommended that it be removed during bathing. Constant swimming and bathing will remove insecticide from the coat and may cause diminished flea and tick control.

TICK CONTROL

The Kiltix Collar will control brown dog ticks and bush ticks. It will provide a high level of control of paralysis ticks but as one female tick is sufficient to harm a dog, the collar should not be relied upon to fully protect the dog. This product is not fully effective against paralysis tick. In paralysis tick areas daily searching of dogs and removal of any ticks found is recommended. In dogs with very thick double coats it may take up to 1 week to establish effective levels in heavily coated areas.

FLEA CONTROL

The Kiltix Collar will start controlling fleas as soon as it is placed around the dog's neck and will continue to do so for 5 months. Fleas are found not only on the dog, but in the dog's surroundings; in the bedding, in the carpet, on the furniture, and in the yard. In order to prevent further infestation of the dog it is important to treat flea larvae in the pet's environment using an appropriate product. Resistance of fleas to propoxur can occur and may result in loss of control.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

SAFETY DIRECTIONS

Poisonous if swallowed. Do not open inner pouch until ready to use. Do not allow children to play with collar. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

DISPOSAL

Dispose of used collars and packaging by wrapping in paper and placing in garbage.

STORAGE

Store below 30°C (Room Temperature).
APVMA Approval No: 51952/59123

Limitation of Liability: Failure to use the product strictly as directed may be illegal, prove dangerous or render the product ineffective. Notwithstanding anything else contained on this label, and subject to any rights you have under the Australian Consumer Law and any other applicable law, Bayer Australia Limited's liability is limited to replacing this product or supplying equivalent product or paying the cost of replacing this product, at its discretion.

KILTIX® is a registered trademark of Bayer AG, Leverkusen, Germany.
Manufactured in Germany.

Bayer

BAYER AUSTRALIA LIMITED
875 Pacific Highway
PYMBLE N.S.W. 2073
ACN 000 138 714

CUSTOMER INFORMATION LINE
1800 678 368 TOLL FREE
From anywhere in Australia 9.00am to 4.00pm (Eastern Time) Mon-Fri
www.friendsforever.com.au



Tick Collar for dogs

kiltix®