

MAIN PANEL – 500 mL**CAUTION**

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**



Solfac[®] Pro

RESIDUAL INSECTICIDE

ACTIVE CONSTITUENT: 25 g/L BETACYFLUTHRIN

GROUP	3A	INSECTICIDE
--------------	-----------	--------------------

**Contents
500 mL**

For the control of cockroaches, spiders, ants, fleas, flies, silverfish, bedbugs, mosquitoes and crickets in the situations specified in the directions for use.

IMPORTANT: Read the attached booklet before use.

APVMA Approval Number: 59424/500mL/0205

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Solfac[®] Pro

RESIDUAL INSECTICIDE

ACTIVE CONSTITUENT: 25 g/L BETACYFLUTHRIN

GROUP 3A

INSECTICIDE

**Contents
200 mL**

For the control of cockroaches, spiders, ants, fleas, flies, silverfish, bedbugs, mosquitoes and crickets in the situations specified in the directions for use.

IMPORTANT: Read the attached booklet before use.

APVMA Approval Number: 59424/200mL/0205

MAIN PANEL – 100 mL**CAUTION****KEEP OUT OF REACH OF CHILDREN****READ SAFETY DIRECTIONS BEFORE OPENING OR USING****Solfac[®] Pro****RESIDUAL INSECTICIDE****ACTIVE CONSTITUENT: 25 g/L BETACYFLUTHRIN****GROUP 3A****INSECTICIDE****Contents
100 mL**

For the control of cockroaches, spiders, ants, fleas, flies, silverfish, bedbugs, mosquitoes and crickets in the situations specified in the directions for use.

IMPORTANT: Read the attached booklet before use.

APVMA Approval Number: 59424/100mL/0205

CAUTION
Keep out reach of children
Read Safety Directions before opening or using

SOLFAC PRO RESIDUAL INSECTICIDE

ACTIVE CONSTITUENT: 25 g/L Betacyfluthrin

GROUP	3A	INSECTICIDE
--------------	-----------	--------------------

For the control of cockroaches, spiders, ants, fleas, flies, silverfish, bedbugs, mosquitoes and crickets in the situations specified in the directions for use.

STORAGE AND DISPOSAL: Store in tightly closed, original container in a safe, well ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. The method of disposal depends on the container type, refer to the disposal statement on the label attached to the container for further directions.

SAFETY DIRECTIONS

When using the prepared spray wear cotton overalls buttoned to the neck and wrist (worn over normal clothing) and a washable hat and elbow length PVC gloves. After use and before eating, drinking or smoking wash hands, arms and face with soap and water. After each days use wash gloves and contaminated clothing.

RE-ENTRY

Do not enter treated areas until the spray has dried.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone 131126). Remove from contaminated area. Apply artificial respiration if not breathing.

MATERIAL SAFETY DATA SHEET

Additional Information is listed in the Material Safety Data Sheet.

Exclusion of Liability

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

SOLFAC® is a registered trademark of Bayer.

Bayer Environmental Science
A Business Group of Bayer CropScience Pty Ltd
A.B.N. 87 000 226 022
391-393 Tooronga Rd
East Hawthorn, Vic. 3123
Phone: (03) 9248 6888
Fax: (03) 9248 6800
Website: www.bayercropscience.com.au

Batch Number: DOM:

For 24 hour specialist advice in an emergency only phone 1800 033 111

Directions for use:**RESTRAINT**

DO NOT apply spray treatment outdoors if rain is imminent or if area is about to be irrigated

Situations	Rate	Pest	Critical Comments
Meatworks (non-product areas), farm buildings, food processing establishments, shops, offices, factories, storerooms, hospitals, schools, barracks, houses, ships, rubbish tips	10 mL/L water		For use on non-porous surfaces; apply at a rate of 5 L of prepared spray per 100 m ² not exceeding the point of run-off. For use on porous surfaces or for use through power equipment; apply at a rate of 10 L per 100 m ² of surface, not exceeding the point of run-off.
		Cockroaches	Thoroughly spray hiding places (eg. cracks and crevices, behind stoves and sinks, in cupboards, around drains, rubbish bins, etc.) until point of run-off. If hiding places are not directly accessible spray barrier strips around them. Repeat treatment as required.
		Web-spinning spiders	Spray spiders, their webbing and areas where spiders may hide directly. It is recommended that webbing is left in place for a few days after application to optimise efficacy.
		Non-web-spinning spiders	Spray areas where spiders are known to occur.
		Ants	Where possible locate the nest and spray directly into nest entrances. Penetration of insecticide can be assisted by pre-wetting sandy and porous soils with water. A thorough soaking treatment with insecticide mixture is required to destroy the nest. If nest cannot be located then spray infested areas.
		Fleas	Clean or vacuum infested areas and burn all dust collected. Lightly spray all infested areas especially under furniture. Pets should be treated with a suitable flea control product registered for that purpose. Do not apply more than 100 mL/m ² to carpet surfaces.
		Mosquitoes and Flies	Spray surfaces where flies and mosquitoes are known to come to rest. Flies – include external surfaces that are exposed to the morning sun. Mosquitoes – spray walls, awnings, fences and vegetated areas in proximity to buildings. Note: The perception of the level of control of flying insects achieved within any given area treated with a residual surface spray will be affected by the extent of in-migration of individuals from external (un-treated) areas. A residual surface spray will only control those insects that alight on the treated surfaces and therefore complete elimination of flying insects from the treated area should not necessarily always be expected..

Directions For Use (Contd):

Situations	Rate	Pest	Critical Comments
Meatworks (non-product areas), farm buildings, food processing establishments, shops, offices, factories, storerooms, hospitals, schools, barracks, houses, ships, rubbish tips	10 mL/L water	Silverfish	Direct spray to areas of potential and existing infestations eg under skirting boards and furniture where dark conditions may favour silverfish populations. If spraying inside cupboards or storage areas remove clothing and utensils etc before spraying.
		Crickets	Spray infested areas. In addition a barrier treatment can be sprayed across areas where these pests enter buildings.
		Bedbugs	Apply to the bed-frame and walls, cracks, crevices and skirting boards in the vicinity of the bed.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

GENERAL INSTRUCTIONS

Mixing: Add the required quantity of Solfac Pro Residual Insecticide to water while stirring.

Application

Do not spray the moving parts of any machinery, electric motors or switchgear with the water-based product.

Note on application to carpets:

Since carpets vary in their thickness, colour and material it is recommended that a small, inconspicuous area of carpet be sprayed and allowed to dry first to ensure that staining will not occur.

Allow treated carpets to dry before resuming activity on them.

GENERAL PRECAUTIONS:

- For surface treatment only – do not spray into the atmosphere using foggers or other devices designed to create an aerosol.
- Use in food processing areas is limited to crack, crevice and barrier strip treatment only.
- When using in animal houses do not spray directly onto livestock, their food, water, litter, feed bins, or surfaces that are accessible to the stock.
- Do not spray directly on humans, exposed food or food utensils or food preparation areas.
- Avoid contact of spray with timber to which a stain is subsequently to be applied. Evidence of water contact on timber is sometimes only evident once the stain has been applied.
- BEFORE USE : Remove or cover all exposed foodstuffs. Cover all dishes and utensils and surfaces where food is prepared.
- AFTER USE : Thoroughly ventilate treated areas. Clean up thoroughly before processing/serving resumes.

INSECTICIDE RESISTANCE WARNING

GROUP	3A	INSECTICIDE
-------	----	-------------

For insecticide resistance management Solfac Pro is a Group 3A insecticide. Some naturally occurring insect biotypes resistant to Solfac Pro and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Solfac Pro or other Group 3A insecticides are used repeatedly. The effectiveness of Solfac Pro on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from failure of Solfac Pro to control resistant insects.

For further information on managing insecticide resistance contact your local supplier or Bayer Environmental Science representative.

Protection of Wildlife, Fish, Crustacea and Environment: Dangerous to fish. Cover fish tanks before using. Do not contaminate ponds, waterways or drains with this product or used container.

Rear panel attached to container

STORAGE AND DISPOSAL: Store in tightly closed, original container in a safe, well ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight.

Rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. Dispose of at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

When using the prepared spray wear cotton overalls buttoned to the neck and wrist (worn over normal clothing) and a washable hat and elbow length PVC gloves. After use and before eating, drinking or smoking wash hands, arms and face with soap and water. After each days use wash gloves and contaminated clothing.

RE-ENTRY

Do not enter treated areas until the spray has dried.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone 131126).

Remove from contaminated area. Apply artificial respiration if not breathing.

MATERIAL SAFETY DATA SHEET

Additional Information is listed in the Material Safety Data Sheet.

Exclusion of Liability

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE

SOLFAC® is a registered trademark of Bayer.

Bayer Environmental Science
A Business Group of Bayer CropScience Pty Ltd
A.B.N. 87 000 226 022
391-393 Tooronga Rd
East Hawthorn, Vic. 3123
Phone: (03) 9248 6888
Fax: (03) 9248 6800

Batch Number: DOM:

For 24 hour specialist advice in emergency only phone 1800 033 111